


MINIMAL REALISTIC $SU(5)$ SCENARIO

Ilja Doršner

*The Abdus Salam International Centre for Theoretical Physics
Strada Costiera 11, 34014 Trieste, Italy*



We present phenomenological aspects of the simplest realistic $SU(5)$ grand unified theory—the theory with the **5**, **15**, and **24** dimensional representations in the Higgs sector. We show that a successful gauge coupling unification sets experimentally accessible upper bound on the total proton decay lifetime. It also relates proton decay lifetime to scalar leptoquark mass in an experimentally testable manner. We also discuss an addition of gauge singlets—both fermions and bosons—to the simplest scenario and comment on relevant phenomenological consequences of such modifications.

1 Introduction

Grand unified theories (GUTs) are considered to be the most viable candidates for the physics beyond the Standard Model. Through matter unification and unification of strong and electroweak interactions they always generate two predictions: (1) gauge couplings unify and (2) proton decays. Testing the first prediction is practically impossible since unification takes place at energy scales beyond our reach. However, the second one can be experimentally investigated thereby offering a very promising way to test grand unification. It is thus important to single out and investigate grand unified theories where proton decay is both accurately predicted and experimentally reachable.

We have recently proposed^{1,2} the simplest realistic GUT model with both of those properties. It comes in a form of a particularly simple extension of the well-known Georgi-Glashow $SU(5)$ model³. In particular, the scenario contains not only the usual three generations of matter fields and the **5** and **24** dimensional Higgs representations, but also one **15** dimensional Higgs. And, it includes *all* possible $SU(5)$ invariant operators.

Despite a large number of parameters in the Lagrangian the scenario is still very predictive. This predictivity is primarily attributed to the simplicity of the Higgs sector. But, two generic features of non-supersymmetric $SU(5)$ framework also boost predictivity. Firstly, $SU(5)$ is

the only simple group with the Standard Model (SM) embedding that has unique single step symmetry breaking. This allows for an accurate determination of the unified scale—the so-called GUT scale M_{GUT} . Secondly, the least model dependent and usually dominant contribution to proton decay comes from an exchange of only one set of superheavy gauge bosons with mass M_V . Clearly, if M_V is taken to define the GUT scale the former property also implies accurate prediction for proton decay. That is exactly what happens in our case.

In the following, we briefly present the simplest realistic $SU(5)$ theory. Again, even though the proposed scenario has uncorrelated regions in the Yukawa sector, the simplicity of its Higgs sector guarantees both its testability and refutability in near future.

2 The simplest $SU(5)$ scenario

The scenario we propose has a following particle content of the Higgs sector¹: $\mathbf{5} = \Psi = (\Psi_D, \Psi_T) = (\mathbf{1}, \mathbf{2}, 1/2) + (\mathbf{3}, \mathbf{1}, -1/3)$, $\mathbf{15} = \Phi = (\Phi_a, \Phi_b, \Phi_c) = (\mathbf{1}, \mathbf{3}, 1) + (\mathbf{3}, \mathbf{2}, 1/6) + (\mathbf{6}, \mathbf{1}, -2/3)$ and $\mathbf{24} = \Sigma = (\Sigma_8, \Sigma_3, \Sigma_{(3,2)}, \Sigma_{(\bar{3},2)}, \Sigma_{24}) = (\mathbf{8}, \mathbf{1}, 0) + (\mathbf{1}, \mathbf{3}, 0) + (\mathbf{3}, \mathbf{2}, -5/6) + (\mathbf{\bar{3}}, \mathbf{2}, 5/6) + (\mathbf{1}, \mathbf{1}, 0)$, where we use the SM ($SU(3) \times SU(2) \times U(1)$) decomposition to set our notation. $\Sigma_{(3,2)}$ and $\Sigma_{(\bar{3},2)}$ are fields eaten by the superheavy gauge fields V . We define the GUT scale through their common mass, i.e., we set $M_V = M_{GUT}$. As always, a vacuum expectation value (VEV) of Σ_{24} breaks $SU(5)$ while VEV of Ψ_D triggers the SM symmetry breaking. During the latter stage Φ_a develops an induced VEV that eventually yields neutrino mass through the type II see-saw mechanism⁴.

On renormalizable level our scenario predicts $Y_D = Y_E^T$ at the GUT scale, where $Y_{D(E)}$ is the down quark (charged lepton) Yukawa coupling matrix. This prediction however disagrees with experiments, especially in the case of the first and second generation. In order to correct that we include higher-dimensional $SU(5)$ invariant operators⁵. (If we demand renormalizability we must introduce $\mathbf{45}$ dimensional Higgs representation⁶. However, falsifiability of such an extension is not guaranteed⁷.)

Clearly, our scenario accommodates realistic fermionic mass spectrum. What remains to be investigated is the issue of gauge unification and its compatibility with proton decay constraints.

2.1 Unification vs. proton decay

There are four masses— M_{GUT} , M_{Σ_3} , M_{Φ_a} and M_{Φ_b} —and two equations that govern gauge coupling unification at the one-loop level¹ in our case. Actually, there are three renormalization group equations—one for each gauge coupling of the SM. However, elimination of the unified coupling constant α_{GUT} leaves only two relevant equations¹⁰. These are

$$\frac{B_{23}}{B_{12}} = 0.719 \pm 0.005, \quad \text{and} \quad \ln \frac{M_{GUT}}{M_Z} = \frac{184.9 \pm 0.2}{B_{12}}, \quad (1)$$

where the right-hand sides reflect the latest experimental measurements of the SM parameters¹¹. The left-hand sides depend on particular mass spectrum of the particle content of the theory at hand. More precisely, $B_{ij} = B_i - B_j$, where B_i coefficients are given by:

$$B_i = b_i + \sum_I b_{iI} r_I, \quad r_I = \frac{\ln M_{GUT}/M_I}{\ln M_{GUT}/M_Z}. \quad (2)$$

b_i are the SM coefficients while b_{iI} are the one-loop coefficients of any additional particle I of mass M_I ($M_Z \leq M_I \leq M_{GUT}$). (Recall, for the case of n light Higgs doublet fields $b_1 = 40/10 + n/10$, $b_2 = -20/6 + n/6$ and $b_3 = -7$.) The B_{ij} -coefficient contributions in our scenario are listed in Table 1. Note that the SM case yields $B_{23}/B_{12} = 0.53$. This means that additional particles

Table 1: B_{ij} coefficients.

	Higgsless SM	Ψ_D	Ψ_T	V	Σ_8	Σ_3	Φ_a	Φ_b	Φ_c
B_{23}	$\frac{11}{3}$	$\frac{1}{6}$	$-\frac{1}{6}r_{\Psi_T}$	$-\frac{7}{2}r_V$	$-\frac{1}{2}r_{\Sigma_8}$	$\frac{1}{3}r_{\Sigma_3}$	$\frac{2}{3}r_{\Phi_a}$	$\frac{1}{6}r_{\Phi_b}$	$-\frac{5}{6}r_{\Phi_c}$
B_{12}	$\frac{22}{3}$	$-\frac{1}{15}$	$\frac{1}{15}r_{\Psi_T}$	$-7r_V$	0	$-\frac{1}{3}r_{\Sigma_3}$	$-\frac{1}{15}r_{\Phi_a}$	$-\frac{7}{15}r_{\Phi_b}$	$\frac{8}{15}r_{\Phi_c}$

with intermediate masses M_I are required for successful unification. In our case these particles are clearly Σ_3 , Φ_a and Φ_b .

In this paper we present the outcome of the two-loop level unification analysis in terms of M_{Σ_3} and M_{Φ_a} contours in the M_{GUT} – M_{Φ_b} plane in Fig. 1. Stars represent points that correspond to exact numerical unification while lines represent linear interpolation.




Figure 1: Unification of the gauge couplings at the two-loop level. Stars correspond to exact numerical two-loop unification solutions. There are two sets of lines of constant value. The steeper set is associated with M_{Φ_a} and the other one represents the lines of constant M_{Σ_3} . All masses are in GeV units. The region to the left of the vertical dashed line is excluded by the proton decay experiments.

The sail-like region in Fig. 1 represents the viable parameter space under the assumption that Ψ_T , Σ_8 and Φ_c are at or above the GUT scale. It is bounded from the left and below by experimental limits on M_{Φ_a} and M_{Φ_b} , respectively. The right bound stems from the requirement that $M_{\Sigma_3} \geq M_Z$. Note that Φ_b is a scalar leptoquark and hence very interesting generator of new physics^{1,2}. For example, Large Hadron Collider (LHC) aims to place more stringent lower limits on the mass of Φ_b at 1 TeV.

The region to the left of the vertical thick dashed line in Fig. 1 is excluded by the present limits on the proton decay lifetime. In order to generate this bound we appropriately take the case of maximal flavor suppression of the gauge $d = 6$ proton decay operators⁹ and use $\alpha = 0.015 \text{ GeV}^3$ for the value of nucleon matrix element¹². Experimental limit we take as input reads $\tau_p(p \rightarrow \pi^0 e^+) > 5.0 \times 10^{33} \text{ years}$ ¹¹.

What happens if we relax the $M_{\Psi_T}, M_{\Sigma_8}, M_{\Phi_c} \geq M_{GUT}$ assumption? As one lowers M_{Ψ_T} and M_{Φ_c} the $M_{\Phi_a} = 130 \text{ GeV}$ line in Fig. 1 moves slowly to the left while, at the same time, the $M_{\Sigma_3} = M_Z$ line moves very rapidly in the same direction until the allowed region shrinks to a point. Hence, any scenario in which M_{Ψ_T} or M_{Φ_c} or both are below M_{GUT} would be more exposed to the tests through the proton decay lifetime measurements and accelerator searches than the scenario shown in Fig. 1.

If, on the other hand, one lowers the mass of Σ_8 , the $M_{\Phi_a} = 130 \text{ GeV}$ line moves to the

right more rapidly than the $M_{\Sigma_3} = M_Z$ line until the allowed region becomes a point when M_{Σ_8} reaches M_Z . At that point M_{GUT} is 4.6×10^{14} GeV for $M_{\Sigma_3} = M_Z$, $M_{\Phi_a} = 6.4 \times 10^3$ GeV, $M_{\Phi_b} = 242$ GeV and $\alpha_{GUT}^{-1} = 37.06$. We use these particular values to derive an accurate upper bound on the proton decay lifetime:

$$\tau_p^{(\text{two-loop})} \leq 1.4 \times 10^{36} \text{ years.} \quad (3)$$

This bound follows from a more general model independent inequality⁹:

$$\tau_p \leq 6 \times 10^{39} \alpha_{GUT}^{-2} (M_V/10^{16} \text{ GeV})^4 (0.003 \text{ GeV}^3/\alpha)^2 \text{ years,} \quad (4)$$

which is applicable to any simple group with the SM embedding. (Note, in the case of scenarios with partial gauge coupling unification such as the Pati-Salam¹³ or flipped $SU(5)$ ^{14,15,16} scenario the proton can be stable¹⁷.)

Clearly, an improvement in the proton lifetime measurements by a 4^4 factor is called for to completely rule out this GUT scenario. The situation is actually even more promising; even a mild improvement in the proton lifetime bounds (by a factor of fifteen) would make our scenario incompatible with exact unification unless either Φ_b or Σ_3 resides below 10^3 GeV. (This certainly makes them accessible in accelerator experiments.) The next generation of proton decay experiments aims at improving lower bounds on partial lifetimes by a few orders of magnitude. For instance, the goal of Hyper-Kamiokande is to explore the proton lifetime at least up to $\tau_p/B(p \rightarrow e^+\pi^0) > 10^{35}$ years and $\tau_p/B(p \rightarrow K^+\bar{\nu}) > 10^{34}$ years in about 10 years¹⁸. Thus, our minimal GUT scenario will be tested and/or ruled out at the next generation of proton decay and accelerator experiments.

3 Adding singlets

Let us comment on possible additions of $SU(5)$ singlets—both spin 1/2 and spin 0 particles—to our scenario.

As we argued, our scenario is tailor made to generate neutrino masses through the Type II seesaw mechanism⁴. But, one can also introduce right-handed neutrinos—singlets of $SU(5)$ —and generate additional neutrino mass contribution through the so-called Type I seesaw mechanism²². That sort of addition obviously cannot change our discussion on the gauge coupling unification and related constraints—at least at the one-loop level. Hence, there is no need to modify our previous discussion on proton decay. However, the addition of right-handed neutrinos has potential to accommodate viable mechanism that explains the origin of the baryon asymmetry observed in the universe¹⁹.

Among viable mechanisms to explain primordial matter-antimatter asymmetry, leptogenesis²⁰ has undoubtedly become one of the most compelling scenarios. In our case, leptogenesis could proceed through the decay of the lightest triplet scalar Φ_a . As has been shown the scalar triplet with a mass $M_{\Phi_a} \geq 10^9\text{--}10^{10}$ GeV²¹ constitutes a natural candidate for a successful triplet leptogenesis if our minimal $SU(5)$ scenario is extended with at least one singlet fermion field. Combining the above leptogenesis bounds with the ones shown in Fig. 1, we observe that successful triplet leptogenesis excludes the region of the parameter space where the mass of the triplet M_{Φ_a} is below $10^9\text{--}10^{10}$ GeV. This in turn would imply that leptoquark Φ_b could be light enough ($M_{\Phi_b} < 10^{6\text{--}7}$ GeV) to open the possibility of direct production at the next generation of collider experiments. It should be noted however that in general if both Type I and Type II seesaws are present leptogenesis could proceed in a way that avoids generation of any bounds on the mass spectrum of relevant particles^{23,24}.

Let us also comment on a possible addition of scalar $SU(5)$ singlet fields from point of view of a successful inflationary scenario^{25,26,27}. Namely, in view of the latest three-year WMAP

data¹⁹ there has been a renewed interest in a class of inflationary models based on a quartic Coleman-Weinberg potential where a gauge singlet scalar field plays inflaton role²⁸. It has been recently shown²⁹ that such a class of models can be in good agreement with WMAP data¹⁹. This conclusion also applies to our scenario if one additional $SU(5)$ singlet scalar field is included.

Finally, another $SU(5)$ scalar singlet can also be introduced to address the existence of Dark Matter¹⁹. This approach however would require additional discrete symmetries³⁰ in order to forbid certain couplings.

4 Additional comments

- Higher-dimensional operators help us generate realistic mass spectrum of the SM fermions. And, these same operators also play a role in deriving a correct upper bound on the total proton lifetime. Clearly, there is a prospect to use this connection to establish an upper bound on scale of the ultraviolet completion of our theory. We will present our findings on this issue in near future[?].
- It is often argued that if Σ_{24} breaks $SU(5)$ down to the SM and Ψ_D gets electroweak VEV v then Σ_3 must get an induced VEV at the tree-level³². We find this to be model dependent statement. Namely, if only $SU(5)$ invariance is imposed then one can always arrange that Σ_3 does not get a VEV at the tree-level by fine-tuning.

5 Summary

We have presented phenomenological aspects of the simplest realistic $SU(5)$ model, where the Higgs sector is composed of the **5**, **15** and **24** dimensional representations. The scenario accommodates realistic fermion masses and gauge coupling unification. Moreover, it predicts experimentally accessible proton decay. More specifically, there exists an upper bound on the total proton decay lifetime $\tau_p \leq 1.4 \times 10^{36}$ years. And, the bound correlates with the leptoquark mass. Since the next generation of proton decay experiments is expected to improve current bounds by a few orders of magnitude, our simplest non-supersymmetric $SU(5)$ model will be certainly tested or ruled out. The proposed scenario, if further extended with singlet fermions and bosons, could accommodate leptogenesis mechanism and be brought in agreement with other cosmological observations.

1. I. Dorsner and P. Fileviez Pérez, Nucl. Phys. B **723** (2005) 53.
2. I. Dorsner, P. F. Perez and R. Gonzalez Felipe, arXiv:hep-ph/0512068.
3. H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32** (1974) 438.
4. G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181** (1981) 287. R. N. Mohapatra and G. Senjanović, Phys. Rev. D **23** (1981) 165.
5. J. R. Ellis and M. K. Gaillard, Phys. Lett. B **88** (1979) 315.
6. H. Georgi and C. Jarlskog, Phys. Lett. B **86** (1979) 297.
7. I. Dorsner and P. F. Perez, arXiv:hep-ph/0606062.
8. A. Giveon, L. J. Hall and U. Sarid, Phys. Lett. B **271** (1991) 138.
9. I. Dorsner and P. Fileviez Pérez, Phys. Lett. B **625** (2005) 88.
10. A. Giveon, L. J. Hall and U. Sarid, Phys. Lett. B **271** (1991) 138.
11. S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
12. Y. Aoki [RBC Collaboration], Nucl. Phys. Proc. Suppl. **140** (2005) 405.
13. J. C. Pati and A. Salam, Phys. Rev. D **10** (1974) 275.
14. A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. Lett. **45** (1980) 413.
15. H. Georgi, S. L. Glashow and M. Machacek, Phys. Rev. D **23** (1981) 783.

16. S. M. Barr, Phys. Lett. B **112** (1982) 219.
17. I. Dorsner and P. Fileviez Perez, Phys. Lett. B **606** (2005) 367.
18. K. Nakamura, Int. J. Mod. Phys. A **18** (2003) 4053.
19. D. N. Spergel *et al.*, arXiv:astro-ph/0603449.
20. M. Fukugita and T. Yanagida, Phys. Lett. B **174** (1986) 45.
21. T. Hambye, M. Raidal and A. Strumia, arXiv:hep-ph/0510008.
22. P. Minkowski, Phys. Lett. B **67** (1977) 421 ; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, eds. O. Sawada et al., (KEK Report 79-18, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen et al., (North-Holland, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, Cargèse, eds. M. Lévy et al., (Plenum, 1980), p. 707; R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44** (1980) 912.
23. T. Hambye and G. Senjanovic, Phys. Lett. B **582** (2004) 73 [arXiv:hep-ph/0307237].
24. T. Hambye, private communication.
25. A. H. Guth, Phys. Rev. D **23** (1981) 347.
26. A. D. Linde, Phys. Lett. B **108** (1982) 389.
27. A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48** (1982) 1220.
28. Q. Shafi and A. Vilenkin, Phys. Rev. Lett. **52** (1984) 691.
29. Q. Shafi and V. N. Senoguz, Phys. Rev. D **73** (2006) 127301.
30. J. McDonald, Phys. Rev. D **50** (1994) 3637.
31. I. Dorsner, P. Fileviez Pérez and G. Rodrigo, in preparation.
32. A. J. Buras, J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, Nucl. Phys. B **135** (1978) 66.